

**MORPHOLOGICAL AND SOME BIOLOGICAL ASPECTS OF THE
PEACH FRUIT FLY, *BACTROCERA ZONATA* (SAUND.)
(DIPTERA: TEPHRITIDAE)**

**HOSNI, MARWA E.¹, M. M. EL-HUSSEINI², A. H. EL-HENEIDY¹
and FATMA A. ATALLAH¹**

1. Dept. of Biological Control, Plant Protection Research Institute, ARC, Giza, Egypt.
2. Dept. of Economic Entomology and Pesticides, Faculty of Agriculture, Cairo University, Giza, Egypt.

(Manuscript received 27 May 2012)

Abstract

The peach fruit fly (PFF), *Bactrocera zonata* (Saunders) (Diptera: Tephritidae) is one of the serious tephritid insect pests attacking tropical and subtropical fruits. In Egypt, the pest is widespread over most of the Egyptian governorates causing serious problems to many fruit crops. The present study aimed to focus on some morphological and biological characteristics of different stages of the PFF under laboratory conditions. Different stages of the PFF were described, measured and illustrated. Biological studies included, durations of different immature stages, fecundity, adult longevity, survival rate and sex ratio. All studies were carried out under the laboratory conditions of $25 \pm 2^{\circ}\text{C}$, 54- 65% R.H. and 14:10 L:D photoperiod. Total developmental period of PFF (from egg deposition to adult emergence) averaged 18.7 days. The female required 13-21days post emergence to start laying eggs. Total number of eggs/female averaged 235.72. Percentage of adult emergence was 58% and sex ratio (females: males) was 1.2: 1.0. Female and male longevity attained 50.6 and 47.3 days for the fed individuals and 2.8 and 3.1 days for unfed ones, respectively.

INTRODUCTION

The dipteran family Tephritidae includes of over 4000 species, of which nearly 700 species belong to Dacine fruit flies (Fletcher, 1987). Nearly 250 species are of economic importance, and distributed widely in temperate, sub-tropical and tropical regions of the world (Christenson and Foote, 1960).

The peach fruit fly (PFF), *Bactrocera zonata* (Saunders) (Diptera: Tephritidae) is one of the serious insect pests attacking tropical and subtropical fruits (Fletcher, 1987) not only in Egypt but also in many other countries. This pest is originally an Indian species, firstly reported from Bengal. It is now widespread over India, Pakistan, Nepal, Bangladesh, Srilanka, Burma and evidently over southeast Asia. In Egypt, the pest is present and widespread over most of the Egyptian governorates causing serious problems to many fruit crops. *B. zonata* has caused an estimation of 190 million EURO damage a year in Egypt (OEPP/EPPO, 2005).

B. zonata has been recorded on over 50 cultivated and wild plant species, mainly those with fleshy fruits. Its main host plants are guava, mango and peach, while the secondary hosts include apricot, fig and citrus. It has been frequently recovered from peach, hence, it is called the peach fruit fly and also due to its serious damage to guava, it is called "guava fruit fly" (Hussain, 1995).

In the last decade, many ecological and biological aspects of the pest were studied to reach proper methods for its control. In Egypt, most researchers initially reared PFF on natural hosts specially, guava fruits to study its biology (Mohamed, 2000). Afterwards, artificial diet was used to obtain large production form eggs and larvae. In 2003, El-Gendy reared larvae of *B. zonata* on wheat bran artificial diet and different host plants and compared the durations of different stages and larval mortality. He reported that the shortest larval duration was recorded on guava while the longest occurred on sour orange in comparison to artificial diet. Amin (2003) evaluated two larval diets, the first one which was reported by Qureshi *et. al.*, (1974), while the other was mentioned by Awadallah (1978) depending upon durations and survival percentages of immature stages. The second diet was most preferable for rearing larvae as it indicated lowest larval duration and highest percentage of larval survival. In addition, it was economic and practical for laboratory manipulation.

El-Naggar (2004) recorded longest incubation period and high hatchability percentages on apple, the shortest was on peach. The longest pupal duration and mortality percentages of *B. zonata* were on apple, peach, guava and apricot, respectively. The PFF was reared at different constant temperatures (15 - 40 °C and 60-70% R.H) by Afia (2007), when incubation periods, larval duration, lower threshold of larval development and average thermal units required under different temperatures were studied. Farag (2009) studied some biological aspects of PFF different stages under three constant temperatures (20, 25 and 30±1°C) to estimate heat-unit requirements necessary for the development of different stages to complete one generation.

Study of insect development and morphological description is a useful tool for evaluation of control programs. Morphology, number of instars, durations of stages is basic aspects of the biology under known conditions that may provide material for comparative studies when the parameters are changed. These results may also be used to develop ecological control models under laboratory or field conditions.

The present study was carried out to shed light on some morphological and biological characteristics of different stages of *B. zonata* under laboratory conditions.

MATERIALS AND METHODS

1. Rearing of *B. zonata*

Flies of *B. zonata* were obtained from a laboratory stock culture at the Biological Control Department, Plant Protection Research Institute, Agricultural Research Center at Giza. Rearing technique was more or less similar to that of the Mediterranean fruit fly, *Ceratitis capitata* (Wied.) but with some variations in the ingredients of the artificial larval diet. The rearing was carried out under the laboratory conditions of 25 ± 2 °C., 54 - 65% R.H. and 14: 10 L: D photoperiod. Eggs of the PFF were collected daily from special egg-laying collector and placed on an artificial diet in plastic trays covered with cloth lids, placed in an incubator at 25 ± 1 °C and left for hatching and larval development. Third larval instars were transferred to trays furnished with a thin layer of sand for pupation. Pupae were sieved from the sand and transferred to adult cages till emergence. Adult flies were reared in cages (35 x 30 x 30 cm).

Flies were encouraged to lay eggs by providing them with egg collectors of plastic mandarins, perforated from the upper part and contained water inside to keep moisture for eggs. Eggs were collected daily and reared on larval artificial diet, which was the same for the Medfly, artificial diet (described by Boller, 1985), but with some modifications as follows: (1000 g short wheat, 250 g molasses or (300 g sugar), 250 g yeast, 10 g sodium benzoate, 1ml Conc. HCL (or 20 ml HCL 2N) and 300 ml water).

Adult flies were fed mainly on sources of sugar, protein and water. Cotton wool soaked with nutritional material served as food source. For protein requirement, mixture of sugar and protein hydrolyzed enzymatic (at a ratio of 3:1, respectively), besides an artificial diet was provided. This diet was described by Masood *et al.* (2006), as follows: (2 pieces Bananas, 6 pieces Egg yolk, 4 table-spoon Honey, 2 table-spoon Vit. B. Complex, 1 table-spoon Yeast, and 8 table-spoon Sugar). The ingredients were mixed in a blender to make a thick syrup solution. For further use, the diet was kept in the refrigerator.

2. Morphological and Biological studies

2.1. Morphology of *B. zonata* immature stages

A set of 50 individuals from each stage was collected from the laboratory stock culture. Each stage was measured and described using a stereomicroscope. For other characteristics of the body such as: anterior, posterior spiracles and mandibles, the larvae were dissected in a saline solution, mounted in Hoyer's medium and described using a microscope (10 x 40).

2.2. Biology of *B. zonata*

2.2.1. Durations of immature stages

Thirty newly deposited eggs were collected from the rearing colony and used as starter for the biological studies. Each individual egg was placed on artificial medium in a small Petri dish and covered with transparent organdy cloth. Petri dishes were incubated at $25 \pm 1^\circ\text{C}$. and eggs were observed daily until hatching. Larvae were left to complete their development inside the medium until occurred on the surface and jumped out from the Petri dishes to pupate in the sand.

2.2.2. Fecundity

Twenty mated females of PFF were used as replicates until their death. Plastic mandarins (had holes in their sides and provided with water) were placed in small cages containing small pieces of the mixture of sugar and protein hydrolyzate and a source of water served as oviposition site. Daily number of eggs laid by each female was counted. Pre-ovipositional, ovipositional and post-ovipositional periods were estimated.

2.2.3. Survival and Sex ratio

Hundred newly formed pupae were collected and placed into a plastic jar covered with organdy until adult emergence. Survival rate was estimated. Numbers of emerged adults were counted.

2.2.4. Longevity

Newly formed pupae were placed individually into small glass vials until adult emergence. Two groups (30 pairs, males and females each) were left to complete their life span, the 1st group was left unfed while the 2nd was fed by a mixture of sugar and protein hydrolyzate and a source of water. Longevity of each sex was estimated.

All biological studies were carried out under the above mentioned laboratory conditions of $25 \pm 2^\circ\text{C}$, 54- 65% R.H. and 14: 10 L: D photoperiod.

3. Statistical analysis was carried out using the F-Test.

RESULTS AND DISCUSSION

1. Morphological Studies of *B. zonata* immature stages

Measurements of different PFF stages are summarized in table (1) and figures (1-12) illustrated characteristics of different stages.

1.1. Egg stage

Eggs of *B. zonata* are deposited by female ovipositor under the skin of the fruits usually in groups. Eggs are white and appear elliptic in shape tapered at both

ends and have a micropyle at one end (Fig. 1). The egg averaged $1220.6 \pm 8.06 \mu$ ($1200 - 1250 \mu$) long and $260 \pm 6.58 \mu$ ($250 - 300 \mu$) wide.

1.2. Larval stage

Larva of the tephritid flies usually passes through three larval instars, described as follows:

1.2.1. First instar larva (L1)

The first instar larva is creamy-white in color, seems typically a mandibulate type larva, with an orange triangular head to support the mandibular muscles (Fig. 2). It measured $1230 \pm 8.06 \mu$ ($1200 - 1250 \mu$) long and $310 \pm 6.58 \mu$ ($300 - 350 \mu$) wide. Mandibles, moderately sclerotized, with relatively small mouth hooks and each with a large preapical hooked tooth (Fig. 5). They measured 100μ long and 50μ wide. In this stage, anterior spiracles are hardly visible under light microscopy (Fig. 6). This note was recorded in *Dacus tryoni* when Elson-Harris (1988) studied the morphology of immature stages of *D. tryoni* and found that the anterior spiracles of 1st instar was reduced to minute simple open pits, hardly visible under light microscopy but visible under scanning electronic microscopy (SEM). Posterior spiracles, each with 2 slits, are obvious but with lightly sclerotized spiracular plates (Fig. 7).

1.2.2. Second instar larva (L2)

The second instar larva is larger than (L1) (Fig. 3) measured $4140 \pm 14.35 \mu$ ($4100 - 4200 \mu$) and $665 \pm 7.54 \mu$ ($650 - 700 \mu$) in length and width, respectively. It has heavily sclerotized, cephalopharyngeal skeleton, well-developed mouth hooks, characterized with medium light color preapical tooth half the size of apical tooth (Fig. 5). The mandibles measured 150μ long and 100μ wide. Anterior and posterior spiracles are clearly observed in this stage (Fig. 7).

1.2.3. Third instar larva (L3)

The third instar larva is creamy-white and appears as a legless maggot (Fig. 5). It feeds on fruit pulp measuring $8258 \pm 7.5 \mu$ ($8250 - 8300 \mu$) long and $1400 \pm 40.34 \mu$ ($1250 - 1500 \mu$) wide. Mandibles are strong, dark and easy to recognize. Each mandible measures 200μ in length and 150μ in width, heavily sclerotised, partially protruding mouth hooks large and symmetrical with curved tips but without preapical teeth on concaved surface (Fig. 5). Anterior and posterior spiracles are similar to those of the 2nd instar but heavily sclerotized (Figs. 6 and 7).

Mature larvae dropped later to the ground through conspicuous holes in the fruit, for pupation in the soil. When disturbed, the larvae double over and jump. The present results agree with those reported by FAO/IAEA (2000) that larvae may grow to a length of about 7-10 mm within the fruit.

1.3. Pupal stage

The third larval instar of the tephritid flies usually pupates in its last molting skin in soil. At the beginning of this stage, the puparium becomes yellow-white. The puparium turns darker and harder by the time (Fig. 8). The full-developed pupa measured $4770 \pm 8.06 \mu$ ($4750-4800\mu$) long and $2010 \pm 6.58\mu$ ($2000-2050\mu$) wide. Puparium is barrel-shaped, reddish brown (Fig. 9), bearing the third instar larva's anterior and posterior spiracles obvious but non-functioning (Figs. 10 and 11).

1.4. Adult stage

B. zonata adult is about the size of the housefly. It is reddish brown with yellowish abdominal cross bands. Wings are transparent with a small brown spot on its tip (Fig. 13). Antennae are short and less than the vertical length of the head. Males and females are easy to recognize through the posterior end of the abdomen (Fig. 14).

Table 1. Measurements of the different stages of *Bactrocera zonata*, all measurements were done by using a micrometer lens (1 mm, scale = 50 μ)

Stage	Mean		Range	
	Length	Width	Length	Width
Egg	1220.6 ± 8.06	260 ± 6.58	1200 - 1250	250 - 300
Larva 1 st instar	1230 ± 8.06	310 ± 6.58	1200- 1250	300 - 350
2 nd instar	4140 ± 14.35	665 ± 7.54	4100 - 4200	650- 700
3 rd instar	8258 ± 7.5	1400 ± 40.34	8250 - 8300	1250 - 1500
Pupa	4770 ± 8.06	2010 ± 6.58	4750 - 4800	2000 - 2050



Fig. 1. Eggs of *B. zonata*



Fig. 2. First instar larva of *B. zonata*



Fig. 3. Second instar larva



Fig. 4. Third instar larva




Fig. 5. Mandibles of A) 1st instar, B) 2nd instar and C) 3rd instar larvae of *B. zonata*



Fig. 6. Anterior spiracle of 1st, 2nd and 3rd instar larvae of *B. zonata*




Fig. 7. Posterior spiracle of 1st, 2nd and 3rd instar larvae of *B. zonata*




Fig. 8. A) Newly yellow-white formed pupa of *B. zonata*
B) Old yellow-brown pupa.



Fig. 9. Pupa of *B. zonata*



Fig. 10. Anterior end of *B. zonata* pupa



Fig. 11. Posterior end of *B. zonata* pupa (takes the form of posterior spiracle)




Fig. 12. Spot on the wing tip of *B. zonata*



Fig. 13. Female and male posterior end of the abdomen of *B. zonata*

2. Biological Studies of *B. zonata*

2.1. Durations of immature stages

The durations of various immature stages of *B. zonata* are presented in table (2).

2.1.1. Incubation period of egg stage

The egg incubation period ranged between 1 and 3 days, with an average of 2.7 days. The present results agree with those of FAO/ IAEA (2000), which reported that under favorable conditions, the eggs hatched to larvae within 2 days. This incubation period may be delayed when temperatures is below normal. Hussain (1995) recorded that the *B. zonata* egg incubation ranged 24–43 hrs. Nearly similar results had been reported by many authors such as, Mohamed (2002), Duyck *et al.* (2004), Afia (2007) and Farag (2009). They reported 3.0, 2.0, 2.3 and 2.2 days at 25°C, respectively.

2.1.2. Developmental period of larval stage

The three larval instars occupied 5-7 days, with an average of 5.8 days. This result is in accordance with the findings of Qureshi *et al.*, (1993), Mohamed (2002), Afia (2007) and Farag (2009) who reported 5.8-12.2, 6.1-13.1, 5.4-14.8 and 5.3-12.8 days at 20–30 °C, respectively as ranges of total developmental larval period.

2.1.3. Duration of pupal stage

As shown in table (2), the duration of pupal stage averaged 10.2 days, ranged between 9 and 11 days. This result was similar to that reported by Qureshi *et al.*, (1993), El-Gendy (2003), Afia (2007) and Farag (2009) who recorded 10.3, 10.8, 11.2 and 10.8 days at 25°C, respectively. Mohamed (2000) reported shorter pupal duration (7.7 days) at the same temperature on the natural host guava fruits.

2.1.4. Total developmental period

Newly emerged adults are not sexually matured, thus the flies attained their sexual maturity within a mean of 16.6 days. The total developmental period of *B. zonata* (from egg deposition to adult emergence) averaged 18.7 days (Table 2).

2.2. Ovipositional periods and fecundity

Table (2) shows that the female required 13-21 days after emergence to start laying eggs. Pre-ovipositional period averaged 16.6 ± 1.97 days at 25°C. The daily mean number of eggs reached 10.5 eggs/ female. The total number of eggs/ female was 235.72 ± 0.24 . In agreement with the obtained results, Qureshi *et al.* (1974), in Pakistan reported that the pre-ovipositional period of *B. zonata* was 14.2 days at 27 ± 2 °C., and the female deposited 279 eggs at 25°C. In (1995), Hussain recorded 13 days prior to female oviposition after emergence. FAO/ IAEA action plan (2000), stated that the pre-ovipositional period, included sexual maturation was 8-16 days and therefore 10 to 23 days were needed to the first egg when the time for sexual maturity was included. The female laid an average of 137 eggs in batches.

The results are also nearly in accordance with those recorded by El-Gendy (2003) who found that the pre-ovipositional period was 14.6 and 26.6 days for

females reared on peach and artificial diets, respectively. Female's fecundity varied from 372 to 734 eggs, when reared on banana and mandarin, respectively. The same trend was recorded by Amin (2003), El-Naggar (2004), Afia (2007), Shehata *et al.* (2008), and Farag (2009). On the other hand, El-Minshawy *et al.*, (1999) recorded a pre-ovipositional period of 45-60 days at 25°C.

Most of the female flies ceased egg deposition for 4-6 days before death. This result is in agreement with that recorded by El-Gendy (2003), Afia (2007) and Farag (2009) who reported 8.0 days at 24±3°C, 5.8 and 6.2 days at 25±1°C, respectively. Amin (2008) recorded 16.4 and 17.2 days at 25°C, when females were fed on solid and liquid protein hydrolysate, respectively.

2.3. Survival rate

Data revealed that the average percentage of adult emergence was 58% at 25±1°C and 54-60 % R.H.

Table 2. Some biological characteristics of *Bactrocera zonata* under the laboratory conditions of 25± 1°C and 54%-60 R.H.

Duration (days)	Means ± SE	Range
Egg	2.7 ± 0.15	1-3
Larva	5.8 ± 0.28	5-7
Pupa	10.2 ± 0.13	9-11
Total developmental period	18.7 ± 0.25	15-21
Survival rate %	58% ± 0.41	40 – 80 %
Longevity (days) d		
Female	50.6 ± 1.44	49- 54
Male	47.3 ± 1.4	44 - 49
Un-fed		
Female	2.8 ± 0.38	2-6
Male	3.3 ± 0.46	2-6
Fecundity		
Pre ovipositional period (days)	16.6 ± 1.97	13 - 21
Ovipositional period	29.33 ± 1.15	28 - 32
Post-ovipositional period	4.6 ± 0.28	4 - 6
Daily mean no. (Eggs/ female)	10.5 ± 1.13	5 -16
Total no. eggs/ female	235.72 ± 6.2	227-248
Sex ratio % (female : male)	1.2 : 1.0	1.3: 1.0 – 2 : 1

2.4. Sex ratio

Sex ratio of *B. zonata* (female: males) was 1.2: 1 under the laboratory conditions of 25±1°C and 54-60% R.H.

2.5. Longevity

When *B. zonata* adults were provided by water and a mixture of sugar and protein hydrolysate, they lived longer compared to those of unfed individuals. The female and male longevities were 50.6 and 47.3 days for the fed individuals and 2.8 and 3.3 for unfed ones, respectively at 25±1°C. This result is nearly similar to that recorded by El-Gendy (2003), Afia (2007) and Farag (2009) who reported these periods for male and female as 59.6 & 82.5, 64.2 & 76.8 and 62.1 & 76.2 days at 25±1°C, respectively. However the obtained results differed than those reported by El- Minshawy et al. (1999) who recorded 100.0 and 145.0 days for male and female at 25°C, respectively .

ACKNOWLEDGEMENTS

The authors wish to thank Dr. Yussef A. Mahmoud, Department of Pests and Plant Protection, National Research Center, Dokki, Giza for his helpful advice for rearing *B. zonata* and providing the diet ingredients.

REFERENCES

1. Afia, Y. E. 2007. Comparative studies on the biology and ecology of the two fruit flies, in Egypt *Bactrocera zonata* (Saunders) and *Ceratitis capitata* (Wiedmann). Ph. D. Thesis, Cairo Univ., Egypt, 301 pp.
2. Amin, A. A. 2003. Studies on peach fruit fly, *Bactrocera zonata* (Saund.) and its control in Fayoum Governorate. M. Sc. Thesis, Fac. Agric., Fayoum Univ., Egypt, 128 pp.
3. Amin, A. A. 2008. Ecological and biological studies on the peach and Mediterranean fruit flies in Fayoum Governorate. Ph.D. Thesis, Fac Agric., Fayoum Univ., Egypt, 251 pp.
4. Awadallah, A. 1978. First report of project for control of Mediterranean fruit fly using sterile insect technique at Fayoum Governorate during the period from June 1974 to June 1978. (PPRI, ARC, library). Cited by Amin (2008).
5. Boller, E. F. 1985. *Rhagoletis cerasi* and *Ceratitis capitata*. In: Sing, P., Moore, R.F., editors. Handbook of Insect Rearing. Elsevier, Amsterdam, the Netherlands, 2: 135-144.

6. Christenson, L. D. and R. H. Foote. 1960. Biology of fruit flies. *Ann. Rev. Ent.*, 5: 171 -192.
7. Duyck, P. F., J. F. Strelin and S. Quilici. (2004): Survival and development of different life stages of *Bactrocera zonata* (Diptera: Tephritidae) reared at five constant temperatures compared to other fruit fly species. *Bull. Ent. Res.*, 94:89-93.
8. El-Gendy, I. R. I. 2003. Studies on peach fruit fly, *Bactrocera zonata* (Saund.) at El-Beheira Governorate. M. Sc. Thesis, Fac. Agric., Alex. Univ., Egypt, 133 pp.
9. El-Naggar, H. M. 2004. Studies on some Apple Insect Pests. M. Sc. Thesis, Fac. Agric., Zagazig Univ., Egypt, 124 pp.
10. El-Minshawy, A. M., M. A. Al-Eryan and A. I. Awad. 1999. Biological and morphological studies on the guava fruit fly *Bactrocera zonata* (Diptera: Tephritidae) found recently in Egypt. *Proceeding of 8th Nat. Conf. of pest and diseases of vegetables and fruits in Ismailia, Egypt*, p.71 -81.
11. Elson-Harris, M. M. 1988. Morphology of the immature stages of *Dacus tryon* (Froggatt) (Diptera: Tephritidae). *J. Aust. ent. Soc.*, 27: 91-98
12. FAO/IAEA (2000): Action Plan: Peach Fruit Fly *Bactrocera zonata* (Saunders). Joint FAO/IAEA Division, Vienna (AT).
13. Farag, Marwa M. 2009. Heat requirements as a tool for predicting the annual generation of peach fruit fly, *Bactrocera zonata* (Saunders) (Diptera: Tephritidae). M. Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, 111 pp.
14. Fletcher, B. S. 1987. The biology of Dacine fruit flies. *Ann. Rev. Entomol.*, 32: 115-114.
15. Hussain, T. 1995. Demography and population genetics of *Dacus zonatus* (Saunders). Ph. D. thesis, Fac. Sci., Punjab Lahore Univ., Pakistan, pp212..
16. Masood, K. K., F S. Mohammad and J. Ghulam. 2006. Effect of different extracts of harmal (*Peganum harmala* L.), rhizomes of kuth (*Saussurea lappa* c. b. clarke) and balchar (*Valeriana officianalis* L.) on the settling and growth of peach fruit fly, (*Bactrocera zonata* Saunders). *Pak. Entomol.*, vol. 28, no.1, 2006.
17. Mohamed, A. M. 2000. Effect of constant temperatures on the development of the peach fruit fly, *Bactrocera zonata* (Saunders) (Diptera: Tephritidae). *Assiut J. Agric. Sci.*, 31: 329- 337.
18. Mohamed, A. M. 2002. Seasonal abundance of the peach fruit fly, *Bactrocera zonata* (Saunders) (Diptera: Tephritidae) with relation to prevailing weather factors in Upper Egypt. *Assiut J. Agric. Sci.*, 33(2): 195-207.
19. OEPP/EPPO. 2005. Data sheets of *Bactrocera zonata*. *Bulletin OEPP/ EPPO Bulletin*. 35: 371-373.

20. Qureshi, Z. A., M. Ashraf, A. R. Bughio and S. Hussain. 1974. Rearing, reproductive behavior and gamma sterilization on fruit fly, *Dacus zonatus* (Saunders) (Diptera: Tephritidae). *Entomol. Exp. Appl.*, 17: 504 – 510.
21. Qureshi, Z. A., T. Hussain, J. R. Carey and R. V. Dowell. 1993. Effects of temperature on development of *Bactrocera zonata* (Saunders) (Diptera: Tephritidae). *Pan-Pacific entomologist*, 69(1): 71-76.
22. Shehata, N.F., M. W.F. Younes and Y.A. Mahmoud. 2008. Biological studies on the Peach Fruit Fly, *Bactrocera zonata* (Saunders) in Egypt. *J. Appl. Sci. Res.*, 4(9): 1103-1106.

مورفولوجيا وبعض الخواص البيولوجية لذبابة ثمار الخوخ

***BACTROCERA ZONATA* (SAUND.) (DIPTERA: TEPHRITIDAE)**

مروة حسنى . *منير الحسيني . أحمد الهندي . فاطمة عطا الله

قسم بحوث مكافحة الحبيبية - معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الجيزة - مصر


* قسم الحشرات الاقتصادية والمبيدات - كلية الزراعة - جامعة القاهرة - الجيزة - مصر

تعتبر ذبابة ثمار الخوخ (*Bactrocera zonata* (Saund.) (Diptera: Tephritidae) واحدة من حشرات عائلة تفرينيتيدي الخطيرة التي تهاجم الفاكهة الاستوائية وشبه الاستوائية. تظهر الآفة في مصر وتنتشر في معظم المحافظات المصرية مسببة مشاكل خطيرة لكثير من محاصيل الفاكهة. هدفت الدراسة الحالية إلى التركيز على بعض الصفات المورفولوجية والبيولوجية للأطوار المختلفة للآفة تحت الظروف المعملية. تم وصف وقياس وتصوير مختلف أطوار الآفة. شملت الدراسات البيولوجية مدة أعمار الأطوار غير الكاملة المختلفة، الخصوبة، طول حياة الحشرة الكاملة، معدل البقاء، والنسبة الجنسية. تمت كل هذه الدراسات تحت الظروف المعملية 25 ± 2 °C و 54-65% رطوبة نسبية، و 14:10 إضاءة : إظلام. بلغ متوسط فترة التطور الكلية للآفة (من وضع البيض إلى الحشرة الكاملة) 18.7 يوماً. احتاجت الأنثى إلى 13-21 يوماً بعد خروجها من طور العذراء لتبدأ في وضع البيض (فترة ما قبل وضع البيض). بلغ متوسط عدد البيض الكلي/ أنثى 235.72 بيضة. كانت نسبة خروج الحشرة الكاملة 58%، وكانت النسبة الجنسية 1.2:1 (إناث: ذكور). بلغ طول حياة الحشرة الكاملة من الإناث والذكور 50.6 و 47.3 يوماً في حالة الأفراد التي تم تغذيتها بينما بلغت 2.8 و 3.1 يوماً في الأفراد غير المغذاة، على التوالي .



VOLUME 91, NUMBER 2
2013





المجلد ٩١ - العدد الثاني

٢٠١٣